

ME/CS 133(a): Homework #1

(Due Monday, October 9, 2017)

Problem 1: (10 points) Let \mathcal{F}_1 denote a fixed reference frame in the plane, with orthonormal basis vectors \vec{x}_1 and \vec{y}_1 . Similarly, consider a second reference frame \mathcal{F}_2 with orthonormal basis vectors \vec{x}_2 and \vec{y}_2 . Let $d_{12} = [x \ y]^T$ be the vector pointing from the origin of \mathcal{F}_1 to the origin of \mathcal{F}_2 . Let θ_{12} denote the relative orientation of the two reference frames: θ_{12} is the angle between \vec{x}_1 and \vec{x}_2 (using the right hand rule, or RHR).

Let ${}^2\vec{v} = [{}^2v_x \ {}^2v_y]^T$ denote the coordinates of a point, P , as seen by an observer in \mathcal{F}_2 . In class we developed a formula for the coordinate transformation of P to its representation in \mathcal{F}_1 :

$${}^1\vec{v} = \vec{d}_{12} + R(\theta_{12}) {}^2\vec{v} \quad (1)$$

where $R(\theta_{12})$ is the 2×2 rotation matrix:

$$R(\theta_{12}) = \begin{bmatrix} \cos \theta_{12} & -\sin \theta_{12} \\ \sin \theta_{12} & \cos \theta_{12} \end{bmatrix}$$

For computational purposes, it is sometimes convenient to use different representations of coordinates, vectors, and rotations. For example, consider complex numbers such that if \vec{w} is a 2×1 vector $\vec{w} = [w_1 \ w_2]^T$ in the plane, then $\tilde{w} = w_1 + iw_2$ where i is the complex number such that $i \cdot i = -1$. Show that if ${}^2\tilde{v}$ is the complex representation of ${}^2\vec{v}$, and \tilde{d}_{12} is the complex representation of \vec{d}_{12} , then the complex representation of the coordinate transform in Equation (1) is:

$${}^1\tilde{v} = \tilde{d}_{12} + e^{i\theta_{12}} {}^2\tilde{v}$$

Problem 2: (10 points) Every planar rigid body displacement is *equivalent* to a rotation about a unique point in the plane, known as the pole (see Figure 1).

Let A be a fixed reference frame. A rigid body, L, which has local frame B attached to it, is located relative to reference frame A by $D_1 = (\vec{d}_{01}, R_{01})$. Body L moves to position C, where the displacement to location C, as measured by an observer in frame B, is given by $D_2 = (\vec{d}_{12}, R_{12})$. Where is the *pole* of the body displacement from position B to position C, as a function of R_{01} , R_{12} , \vec{d}_{01} , and \vec{d}_{12} ?

- As measured in Frame A
- As measured in Frame B
- As measured in Frame C

Problem 3: (5 points) In the above problem, suppose $D_1 = (x, y, \theta) = (1.0, 2.0, 30.0^\circ)$ and $D_2 = (x, y, \theta) = (2.0, 2.0, 45^\circ)$. Where is the pole of the displacement from B to C in this case?




Figure 1: Geometry of planar displacement

Problem 4: (15 points) Using the set up of Problem 2, pick a coordinate system whose origin is located at the pole of the displacement, and show that in this coordinate system, the displacement of the body from B to C is a pure rotation. That is, in that coordinate frame, the displacement should take the form $D = (\text{displacement vector}, \text{rotation matrix}) = (\vec{0}, R)$, where R is a 2×2 rotation matrix.

Problem 5: (15 points) A *planar reflection* is an operation wherein one “reflects” all of the particles in a body across a line (see Figure 2). Show (intuitively) that reflections “preserve length.” That is, reflections do not alter the distance relationship between particles in a rigid body. Can any planar displacement be equivalently performed by a reflection?




Figure 2: Geometry of Planar Rigid Body Reflection

NOTE: This homework is due at 5:00 pm on Oct. 9. You can turn in your homework solution in class (which occurs from 3:00-4:00 pm that day), or leave the solution with Ms. Sonya Lincoln, room 250 of the Gates-Thomas Building